

List of tables

Table 1.1 – Description of main features of the six SEAs	4
Table 1.2 – Interim ecological objectives, and expected outcomes for the six SEAs	6
Table 2.1 – Equivalence of three different vegetation classification schemes applied to the Barmah–Millewa Forest	17
Table 2.2 – Areas of key floodplain plant associations at Barmah and Millewa forests, estimated from aerial photograph, satellite images and field survey	19
Table 2.3 – Flood frequencies of the major Barmah–Millewa Forest floodplain vegetation communities before regulation	20
Table 2.4 – River Murray flows (Tocumwal) at which the main vegetation communities commence to fill	21
Table 2.5 – Altered flooding and drying characteristics of the main vegetation communities in the Barmah Forest.	24
Table 2.6 – Impact of hydrological changes on vegetation composition of Barmah Forest	24
Table 3.1 – Major effluents into Gunbower and Koondrook–Perricoota forests	53
Table 3.2 – Equivalence of different vegetation classification schemes applied to the Gunbower Forest	56
Table 3.3 – Flood frequencies of the major Gunbower Forest floodplain vegetation communities before regulation	60
Table 3.4 – Relationship between river flow, forest inflow and inundated area by Red gum vegetation type in the Koondrook–Perricoota Forest	63
Table 3.5 – Impact of environmental changes on vegetation communities of Gunbower Forest	66
Table 3.6 – Priority of WRCs for water management	70
Table 3.7 – Effects of regulation of the River Murray on flows at Torrumbarry, based on modelled monthly flows	75
Table 3.8 – Effect of river regulation on the frequency of a range of flow events in Spring months over period 1891-2000 in the Koondrook–Perricoota Forest	76
Table 4.1 – Critical flow threshold at Euston for flooding of the 18 Hattah Lakes and Chalka Creek under current and natural conditions	90
Table 4.2 – Estimated holding capacity for the 18 Hattah Lakes	90
Table 4.3 – Registered cultural heritage sites in the Hattah–Kulkyne National Park	106
Table 4.4 – Frequency and type of significant Indigenous sites located in the Hattah–Kulkyne National Park	107
Table 4.5 – Percentage of time a flow event is exceeded using daily data for 109 years for modelled natural and current conditions at Euston Weir	116
Table 4.6 – Percentage of years when an event greater than the threshold flow occurs for at least 14 days duration; using daily data for 109 years, for modelled natural and current conditions at Euston Weir	116
Table 4.7 – Median duration of events (of ≥ 14 days duration) for given flow thresholds using daily data for 109 years for modelled natural and current conditions at Euston Weir	117

Table 4.8 – Level of risk to Ramsar values at the Hattah–Kulkyne Lakes Ramsar site	119
Table 5.1 – Flood frequencies of the major Chowilla Floodplain vegetation communities before regulation	138
Table 5.2 – River Murray flows at which high conservation value floodplain wetlands at Chowilla Floodplain commence to fill, and the change in frequency of those flows	138
Table 5.3 – The effects of River Murray flows on floodplain inundation, flooding frequency, and flood duration under natural and current conditions at Chowilla Floodplain, based on MDBC monthly simulation model analysis for period to 1995	156
Table 5.4 – Predicted percentage of the total existing area of the nine major vegetation communities inundated by the two natural (and enhanced) 2000 flood peaks on Chowilla Floodplain	173
Table 5.5 – Flow thresholds for landscape feature inundation for both the Lindsay and Wallpolla Islands	179
Table 5.6 – Flow thresholds of the River Murray, above which anabranches and floodplains receive inflows for Natural and Current conditions	204
Table 5.7 – Percentage of time a flow event is exceeded using daily data for 109 years for modelled Natural and Current conditions at Lock 10, 9 and 7	206
Table 5.8 – Number of events that flow exceeds a given threshold and for a duration of 14 days or more using daily data for 109 years for modelled natural and current conditions at Lock 10, 9 and 7.	207
Table 5.9 – Median duration of events (of 14 days or more) for given flow thresholds using daily data for 109 years for modelled natural and current conditions at Locks 10, 9 and 7.	208
Table 7.2 – Main functions of each MDBC flow management structure	240
Table 7.3 – Characteristics of weirs and weir pools on the lower River Murray	242
Table 7.4 – Formalised operating procedures to protect environmental values of the River Murray Channel	246
Table 7.5 – Operating practices to protect environmental values of the River Murray Channel	247
Table 7.6 – Major salt interception schemes along the River Murray	257
Table 7.7 – Main processes threatening ecological health along the River Murray Channel	265
Table 7.8 – Definitions of abbreviations and explanation of ecological significance of elements of the generalised flow regime illustrated on Figure 7.22	268
Table 8.1 – Distinctive features of the significant ecological assets	276